

Vortex Waves in a Cloud of Bose Einstein - Condensed, Trapped Alkali - Metal Atoms

Carlo F. Barengi

Mathematics Department
University of Newcastle upon Tyne
NE1 7RU England

Abstract

We consider the vortex state solution for a rotating cloud of trapped, Bose Einstein - condensed alkali atoms and study finite temperature effects. We find that thermally excited vortex waves can distort the vortex state significantly, even at the very low temperatures relevant to the experiments.

PACS numbers 0.35 Fi, 5.30 Jp, 32.80 Pj

Recent experiments [1,2,3] have demonstrated the existence of Bose - Einstein condensation [4,5,6] in clouds of trapped alkali atoms and generated great theoretical interest [7,8]. To understand the superfluid aspects of these new systems it is natural to study what happens when the clouds are rotated [9]. The existence of quantised vortex lines in rotating clouds has not been experimentally established yet. Nevertheless a vortex state solution of the governing Gross - Pitaevskij equation has been found already by Dalfovo and Stringari [10]. They solved the equation for systems of $N \approx 10^4$ atoms at temperature $T = 0$ flowing with quantised circulation around a vortex line set on the z axis of rotation of the cloud. They also argued that the new solution is relevant to the stability [11] of trapped clouds of atoms which have negative scattering length, such as ^7Li [3]. Dalfovo and Stringari's work has extended to new systems the study of quantised vorticity, until now restricted to superfluid helium and neutron stars [12], a development which is important as new non - destructive detection methods are invented [13].

The aim of this brief report is to estimate some finite temperature corrections. We find that the vortex state calculated at $T = 0$ by Dalfovo and Stringari can be rather distorted at small but finite T by sinusoidal displacements of the vortex core away from the axis of rotation. These displacements, called *vortex waves* [14,15], are excited thermally. The effect is at first surprising, given the very low temperatures at which clouds of trapped Bose Einstein - condensed atoms are produced, about $10^{-7}K$ only.

Let us consider a cloud of size $L = 2R_c$ of trapped, Bose Einstein - condensed atoms at temperature T . The cloud rotates about the z axis at angular velocity $\Omega \geq \Omega_{crit}$, where the critical velocity Ω_{crit} for the appearance of the first vortex line is typically of the order of few Hertz [6,10]. The vortex line is located on the z axis in its unperturbed state at $T = 0$. Let us assume that at temperature $T > 0$ the vortex core is displaced away from the axis of rotation by the amount

$$\eta(z, t) = A \cos(\omega t) \cos(kz), \quad (1)$$

where λ , $k = 2\pi/\lambda$, ω and A are respectively the wavelength, the wavenumber, the angular frequency and the amplitude of the wave. In the first

approximation we assume that the cloud is large enough that we can neglect size effects [16] on the dispersion relation $\omega = \omega(k)$ of the wave. We have then $\omega = \Gamma k^2 \mathcal{L}/4\pi$ [12], where $\Gamma = h/m$ is the quantum of circulation, h is Plank's constant, m is the mass, $\mathcal{L} \approx \ln(2/ka_{core})$ and a_{core} is the vortex core parameter. To determine the amplitude of a vortex wave of wavenumber k we note that it is a linear perturbation of the vortex state solution of the Gross - Pitaevskij equation [17], in the same way as phonons are perturbations of the uniform solution. Assuming that vortex waves are quantized like harmonic oscillators, the energy $\mathcal{E}$ of the wave of wavenumber k at temperature T is equal to the energy $\hbar\omega$ times the mean number $1/[\exp(\beta\hbar\omega) - 1]$ of excitations in that state k . There are two contributions to $\mathcal{E}$: the kinetic energy $\mathcal{E}_1$ of the fluid rotating around the core and the potential energy $\mathcal{E}_2$ due to the increase $\delta\ell$ of the vortex' length from the unperturbed to the pertuebed state. We have

$$\frac{\hbar\omega}{e^{\beta\hbar\omega} - 1} = \mathcal{E}_1 + \mathcal{E}_2, \quad (1)$$

where $\beta = 1/K_B T$ and K_B is Boltzmann's constant. The kinetic energy is obtained from averaging the energy density $\rho_{eff}\pi a_{core}^2 \dot{\eta}^2/2$ over the period $2\pi/\omega$ of oscillation and the dimension of the system, where $\rho_{eff} = \rho/2$ is the hydrodynamic mass and a_{core} is the size of the vortex core:

$$\mathcal{E}_1 = \frac{\pi}{8} a_{core}^2 \rho_{eff} A^2 \omega^2 L, \quad (2)$$

The potential energy is $\mathcal{E}_2 = \epsilon\delta\ell$ where ϵ is the energy of the vortex motion per unit length in the z direction, $\ell = L$ is the length of the unperturbed vortex and $\ell' = \int_0^L dz \sqrt{1 + A^2 k^2 \cos^2(kz)}$ is the length of the vortex in the presence of waves. To obtain ϵ we integrate $\rho v^2/2$ around the vortex line, where $v = \Gamma/2\pi r$ is found from the quantization of circulation. We obtain

$$\mathcal{E}_2 = \delta\ell \frac{\rho\Gamma^2}{4\pi} \ln(R_c/a_{core}), \quad (2)$$

The largest amplitudes arise from the smallest wavenumbers, so we assume $\lambda \approx L = 2R_c$. In the limit $Ak \ll 1$ we can Taylor expand the integrand for ℓ' . We also note that typically $\beta\hbar\omega \ll 1$, and, neglecting a term proportional to $a_{core}^2/R_c^2 \ll 1$, we conclude that

$$A \approx \sqrt{\frac{32K_B T R_c}{\pi \rho \Gamma^2 \ln(R_c/a_{core})}}, \quad (4)$$

We apply equation (4) to the ^{87}Rb experiments of Ref. 1 and the discussion of Dalfovo and Stringari [10]. The quantum of circulation is $\Gamma = 4.4 \times 10^{-5} \text{ cm}^2/\text{sec}$. The average density of atoms in typical clouds is estimated to be in the range from $n \approx 10^{12}$ to 10^{13} cm^{-3} which corresponds to $\rho \approx 10^{-10}$ to 10^{-9} g/cm^3 . The vortex core parameter enters equation (4) only via a slow logarithmic term. From the calculation of Dalfovo and Stringari for a small cloud of $N = 5,000$ atoms (their figure 4b) we infer that a_{core} , which we define for convenience as the distance from the axis over which the computed condensate's wave function recovers half of its peak value, is $a_{core} \approx 0.2a_{\perp} \approx 0.24 \times 10^{-4} \text{ cm}$ where $a_{\perp} \approx 1.2 \times 10^{-4} \text{ cm}$ is the oscillator's characteristic length. At $T = 10^{-7} \text{ K}$ and $R_c \approx 5 \times 10^{-4} \text{ cm}$ [6,10] we find that A ranges from 0.33×10^{-3} to $0.11 \times 10^{-3} \text{ cm}$. This amplitude is not a negligible distortion of the vortex state: it represents a displacement of the vortex line of a distance from 4 to 14 times the size of the core. At larger values of R_c we expect size effects to become less important for which equation (4) becomes a better estimate of A . For example at $R_c \approx 10^{-3} \text{ cm}$ the displacements are 6 to 20 times larger than a_{core} . It is instructive to compare this result with the case of a vortex line in superfluid ^4He . In ^4He the vortex waves are negligible. Using $\rho = 0.145 \text{ g/cm}^3$ and $a_{core} \approx 10^{-8} \text{ cm}$ [18] and the same $R_c = 5 \times 10^{-3} \text{ cm}$ and $T = 10^{-7} \text{ K}$ used before, we find $A \approx 0.2 \times 10^{-9} \text{ cm}$, a displacement which is only two hundredths of helium's vortex core size.

Finally we compare the thermally induced displacements of the core with the amplitude of the zero - point motion of the vortex line, $A_0 \approx 0.5\sqrt{2\hbar/\rho\Gamma a_{core}}$ [19]. Using the two values of ρ mentioned above we find that A_0/a_{core} ranges from 1 to 3, which is less than the ratio A/a_{core} for the thermal displacements. For vortex lines in ^4He one has $A_0/a_{core} = 2$.

We conclude that long wavelength thermal waves can affect the vortex state even at the very low temperatures relevant to the experiments.

References

- [1] M.H. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wieman & E.A. Cornell, *Science* **269**, 198 (1995).
- [2] K.B. Davis, M.O. Mewes, N.J. van Druten, D.S. Durfee, D.M. Kurn & W. Ketterle, *Phys. Rev. Letters* **75**, 3969 (1995).
- [3] C.C. Bradley, C.A. Sackett, J.J. Tollett & R.G. Hulet, *Phys. Rev. Lett.* **75**, 1687 (1995).
- [4] A. Griffin, D.W. Snoke & S. Stringari ed's. *Bose Einstein Condensation* Cambridge University Press (1996).
- [5] A.L. Fetter, *Phys. Rev. A* **53**, 4245 (1996).
- [6] G. Baym & C.J. Pethick, *Phys. Rev. Letters* **76**, 6 (1996).
- [7] M. Edwards & K. Burnett, *Phys. Rev. A* **51**, 1382 (1995).
- [8] P.A. Ruprecht, M.J. Holland, K. Burnett & M. Edwards, *Phys. Rev. A* **51**, 4704 (1995).
- [9] S. Stringari, *Phys. Rev. Letters* **76**, 1405 (1996).
- [10] F. Dalfovo & S. Stringari, *Phys. Rev. A* **53**, 2477 (1996).
- [11] L.P. Pitaevskij, preprint cond-mat/9605119 (1996).
- [12] R.J. Donnelly, *Quantized Vortices in Helium II*, Cambridge University Press (1992).
- [13] M.R. Andrews, M.O. Mewes, N.J. Vandruten, D.S. Durfee, D.M. Kurn & W. Ketterle, *Science* **273**, 84 (1996).
- [14] W.I. Glaberson R.J. Donnelly, in *Progress in Low Temperature Physics* vol. IX, edited by D.F. Brewer, Elsevier Science Publishers, page 1 (1986)

- [15] C.F. Barenghi, R.J. Donnelly & W.F. Vinen, Phys. Fluids **28**, 498 (1985).
- [16] In the case of phonons, size effects for small clouds have been calculated by S. Stringari, preprint cond-mat/9603126 (1996) and M. Edwards, P.A. Ruprecht, K. Burnett, R.J. Dodd & C.W. Clark, preprint cond-mat / 9605170 (1996).
- [17] L.P. Pitaevskij, Sov. Phys. JETP **13**, 451 (1961)
- [18] C.F. Barenghi, R.J. Donnelly & W.F. Vinen, J. Low Temp. Phys. **52**, 189 (1982).
- [19] A.L. Fetter, Phys. Rev. **162**, 143 (1967).